

**Practice Problems**

These problems are not to be handed in, but try them first; do as many of them as you need until they're easy, or make up more along the same lines if you need more practice.

**1** Suppose that  $x$  is a variable quantity and suppose that  $y = 3x$ .

- What is  $\Delta_4^5(x^2)$ ?
- What is  $\Delta_{x=-1}^{x=4}(xy)$ ?
- What is  $\Delta_7^9(y^2)$ ?
- What is  $\Delta_5^3(x + 9)$ ?

**2** For each of the following expressions, find its antiderivatives (indefinite integrals).

- $3x^2 dx$
- $(5x^3 - 3x + 4) dx$
- $\left(\sqrt{y} + \frac{1}{y^2}\right) dy$
- $\left(2x + \frac{2}{x}\right) dx$

**3** Evaluate each of the following definite integrals.

- $\int_0^5 3x^2 dx$
- $\int_{-1}^3 (5x^3 - 3x + 4) dx$
- $\int_1^2 \left(\sqrt{y} + \frac{1}{y^2}\right) dy$
- $\int_4^3 \left(2x + \frac{2}{x}\right) dx$

**Due Problems**

These problems are due May 15 Tuesday.

**1** For each of the following expressions, find its antiderivatives (indefinite integrals). Show at least one intermediate step for each.

- $(2x^3 + 4x) dx$
- $\left(\frac{2}{x^3} + \sqrt[3]{x} + \frac{2}{x}\right) dx$

**2** Using your answers to Problem 1 above, evaluate each of the following definite integrals. Show at least one additional intermediate step for each.

a.  $\int_3^4 (2x^3 + 4x) dx$

b.  $\int_{-8}^{-1} \left( \frac{2}{x^3} + \sqrt[3]{x} + \frac{2}{x} \right) dx$

### Answers to Practice Problems

Here are the answers to the Practice Problems from the beginning of the assignment.

**1**

a. Just plug in and subtract:

$$\Delta_4^5(x^2) = ((5)^2) - ((4)^2) = 25 - 16 = 9.$$

b. Since  $y = 3x$ ,

$$\Delta_{x=-1}^{x=4}(xy) = \Delta_{x=-1}^{x=4}(x(3x)) = \Delta_{-1}^4(3x^2) = (3(4)^2) - (3(-1)^2) = 48 - 3 = 45.$$

c. This is not a fair question! It *might* mean

$$\Delta_{y=7}^{y=9}(y^2) = ((9)^2) - ((7)^2) = 81 - 49 = 32,$$

but it *might* mean

$$\Delta_{x=7}^{x=9}(y^2) = \Delta_{x=7}^{x=9}((3x)^2) = \Delta_7^9(9x^2) = (9(9)^2) - (9(7)^2) = 729 - 441 = 288.$$

I should never give you an unclear problem like this one!

d. This is also kind of a trick question, but it is fair, and I might give you a problem like it.

$$\Delta_5^3(x+9) = ((5)+9) - ((3)+9) = 14 - 12 = 2.$$

**2**

a. Running the power rule in reverse, I add 1 to the exponent and divide the coefficient by the new exponent.

$$3x^2 dx = d\left(\frac{3}{3}x^3\right) = d(x^3),$$

so

$$\int 3x^2 dx = x^3 + C,$$

where  $C$  is an arbitrary constant.

b. Now I run the power rule in reverse for each term.

$$(5x^3 - 3x + 4) dx = (5x^3 - 3x^1 + 4x^0) dx = d\left(\frac{5}{4}x^4 - \frac{3}{2}x^2 + \frac{4}{1}x\right) = d\left(\frac{5}{4}x^4 - \frac{3}{2}x^2 + 4x\right),$$

so

$$\int(5x^3 - 3x + 4) dx = \frac{5}{4}x^4 - \frac{3}{2}x^2 + 4x + C,$$

where  $C$  is an arbitrary constant.

c. I rewrite using exponents so that I can reverse the power rule.

$$\left(\sqrt{y} + \frac{1}{y^2}\right) dy = (1y^{1/2} + 1y^{-2}) dy = d\left(\frac{1}{3/2}y^{3/2} + \frac{1}{-1}y^{-1}\right) = d\left(\frac{2}{3}\sqrt{y^3} - \frac{1}{y}\right),$$

so

$$\int\left(\sqrt{y} + \frac{1}{y^2}\right) dy = \frac{2}{3}\sqrt{y^3} - \frac{1}{y} + C,$$

where  $C$  is an arbitrary constant.

d. When the exponent is  $-1$ , the antiderivative involves a logarithm.

$$\left(x + \frac{1}{x}\right) dx = (1x^1 + 1x^{-1}) dx = d\left(\frac{1}{2}x^2 + 1 \ln |x|\right) = d\left(\frac{1}{2}x^2 + \ln |x|\right),$$

so

$$\int\left(x + \frac{1}{x}\right) dx = \frac{1}{2}x^2 + \ln |x| + C,$$

where  $C$  is an arbitrary constant.

### 3

a.

$$\int_0^5 3x^2 dx = \int_0^5 d(x^3) = \Delta_0^5(x^3) = ((5)^3) - ((0)^3) = 125 - 0 = 125.$$

b.

$$\begin{aligned}\int_{-1}^3 (5x^3 - 3x + 4) dx &= \int_{-1}^3 d\left(\frac{5}{4}x^4 - \frac{3}{2}x^2 + 4x\right) = \Delta_{-1}^3\left(\frac{5}{4}x^4 - \frac{3}{2}x^2 + 4x\right) \\ &= \left(\frac{5}{4}(3)^4 - \frac{3}{2}(3)^2 + 4(3)\right) - \left(\frac{5}{4}(-1)^4 - \frac{3}{2}(-1)^2 + 4(-1)\right) \\ &= \frac{399}{4} - \left(-\frac{17}{4}\right) = 104.\end{aligned}$$

c.

$$\begin{aligned}\int_1^2 \left( \sqrt{y} + \frac{1}{y^2} \right) dy &= \int_1^2 d \left( \frac{2}{3} \sqrt{y^3} - \frac{1}{y} \right) = \Delta_1^2 \left( \frac{2}{3} \sqrt{y^3} - \frac{1}{y} \right) \\ &= \left( \frac{2}{3} \sqrt{(2)^3} - \frac{1}{(2)} \right) - \left( \frac{2}{3} \sqrt{(1)^3} - \frac{1}{(1)} \right) \\ &= \left( \frac{4}{3} \sqrt{2} - \frac{1}{2} \right) - \left( -\frac{1}{3} \right) = -\frac{1}{6} + \frac{4}{3} \sqrt{2} \approx 1.72.\end{aligned}$$

d.

$$\begin{aligned}\int_3^4 \left( 2x + \frac{2}{x} \right) dx &= \int_3^4 d \left( \frac{1}{2} x^2 + \ln |x| \right) = \Delta_4^3 \left( \frac{1}{2} x^2 + \ln |x| \right) \\ &= \left( \frac{1}{2} (4)^2 + \ln |4| \right) - \left( \frac{1}{2} (3)^2 + \ln |3| \right) \\ &= (8 + 2 \ln 2) - \left( \frac{9}{2} + \ln 3 \right) = \frac{7}{2} + 2 \ln 2 - \ln 3 \approx 3.79.\end{aligned}$$